



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

"The Isolation of *Bacillus Typhosus* from Butter," D. H. Bergey.

"Note on a Peptid-splitting Enzyme in Woman's Milk," Louis M. Warfield.

"Carcinoma Involving the Entire Kidney" (with two plates), Lindsay S. Milne.

"A Study of a Case of Thrombo-angitis Obliterans," Harlow Brooks.

"The Value of the 'Hormone' Theory of the Causation of New Growth," I. Levin and M. J. Sittenfeld.

THE contents of the *Astrophysical Journal* for September are:

"Spectrum of Comet Morehouse (1908 c)," A. de la Baume Pluvinel and F. Baldet.

"The Discovery of Eclipsing Variable Stars," Joel Stebbins.

"A New Bright Variable Star, β Aurigæ," Joel Stebbins.

"Motion and Condition of Calcium Vapor over Sun-spots and other Special Regions. II.," Charles E. St. John.

"An Enclosed Arc for Spectroscopic Work," James Barnes.

"The Spectra of Aluminium, Copper and Magnesium in the Arc under Reduced Pressure," James Barnes.

"An Inquiry into the Variation of the Spectroscopic Binary κ Pavonis," Alex. W. Roberts.

SPECIAL ARTICLES

THE ORIGIN OF THE GREAT PLAINS

PASSARGE's dictum that "Wasser ist nicht im Stande solche Ebene zu erodieren" now seems eminently applicable to vast, arid and remarkably smooth plains other than those of the great South African plateau. In the light of the recent advances in our knowledge of general desert-leveling, or regional planation and lowering without base-leveling, the vastness and evenness of the Great Plains lying between the Rocky Mountains and Mississippi River at once raise the query whether genetically their dominant characters have been properly interpreted.

At the present time the geologic formations receiving greatest critical attention are those known as continental deposits, or terranes laid down and preserved on land instead of in lakes or seas. In the recent considerations of

the subaerial formations so many novelties enter that in many an old and well-known field a new interest is aroused. The Great Plains and their deposits are one of these. On a grand scale they appear to introduce to us a mode of terranal genesis hitherto almost unrecognized. Continental deposits thus begin to assume in this country an importance which has never been before accorded them.

Singularly enough, the so-called fresh-water Tertiaries of the Great Plains have had ascribed to them every known method of origin. The same is true of the surface-relief. In the descriptions and discussions of this one geologic formation and of this single topographic feature is reflected in all its various phases a century's trend of sedimentative and physiographic thought in America. For this reason, if for no other, the theme is deserving of more than passing notice.

The origin of the Great Plains and their deposits has been ascribed to (1) normal marine deposition, (2) lacustrine sedimentation in vast bodies and (3) fluvial aggradation. To these hypotheses must now be added a fourth—that of eolian planation. In the extreme west in front of the Rocky Mountains is a belt of deflative character where often the substructure forms a typical rock-floor. In the broad median belt eolian deposition to vast extent has taken place, and is still going on. In the eastern belt along the Missouri River wind-effects, although extensive, are almost wholly obscured by moist-climate phenomena.

The two essential points to be noted are, first, continental deposits may be as important as marine or lacustrine deposits; and second, that on the American continent eolic deposits are of vast extent and are being formed under conditions whereby they may be preserved through the geologic ages as effectually as any of the marine Cambrian terranes have been.

In this new century the theory of eolic planation and deposition promises to be one of the half-dozen great and novel thoughts in the domains of geology.

CHARLES R. KEYES